

K-Rich™

Problems with potassium uptake are often pH-related, due to the inherent alkalinity of this element. K-Rich™ involves potassium citrate – a buffered, super-available form of potassium.

High analysis pH-buffered liquid potassium complexed with citric acid. K-Rich™ is specifically designed for foliar fertilising fruit, vegetable and grain crops during the potassium-hungry fruit/seed-fill stage.

BENEFITS

- Australian Certified Organic (ACO) Registered Farm Input 456AI.
- Super available, high analysis potassium essential for optimum flower/fruit/seed production.
- Clear formulation ideal for all crop stages.

PACKAGING:

1 L, 5 L, 20 L,
200 L, 1000 L

Product Code: KRICH



**PREMIUM
LIQUIDS**



APPLICATION RATES

Foliar Spray:

Vegetables

1 L per 100 L water up to 5 L/ha equivalent
Apply weekly from fruit-set until harvest or as required

Orchards & Vineyards

1 L per 100 L water up to 7 L/ha equivalent
Apply every 2 – 4 weeks from fruit-set until harvest or as required

Ornamentals

1 L per 100 L water up to 5 L/ha equivalent
Apply every 2 – 4 weeks from fruit-set until harvest or as required

Turf

1 L per 100 L water up to 5 L/ha equivalent
Apply every 2 – 4 weeks or as required

Broadacre Crops

2 – 3 L/ha in 60 – 100 L water, depending on canopy closure
Apply as required

Pasture

2 – 3 L/ha in 60 – 100 L water
Apply as required

Spot-Spraying

50 mL per 10 L water
Apply as required

Home Garden Application:

Dilute 30 mL per 9 L watering can and apply the diluted mix at a rate of 1 litre per square metre
Apply fortnightly or as required

TYPICAL ANALYSIS W/V

Potassium (K) (as citrate)	33.29%
SG	1.5
pH	7 – 8
Conductivity	20 – 40 mS/cm
Appearance	Clear liquid

INSTRUCTIONS

Shake or stir well before use. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application. Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application. Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use.

IMPORTANT

- **Do not spray if copper hydroxide residue may be present.**
- **Note:** Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

STORAGE & HANDLING

- Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 15°C. Crystallisation or sedimentation may occur below 15°C. Do not pre-mix or store in diluted form. Seal lid/cap immediately after use.

Disclaimer: Any recommendations provided by Nutri-Tech Solutions Pty Ltd (NTS) or its Distributors are advice only. As no control can be exercised over storage, handling, mixing, application or use, or weather, plant or soil conditions before, during or after application (all of which may affect the performance of our program), no responsibility for, or liability for any failure in performance, losses, damages, or injuries (consequential or otherwise), arising from such storage, mixing, application, or use will be accepted under any circumstances whatsoever. NTS recommend you contact an Agronomist prior to product application. The Buyer assumes all responsibility for the use of any NTS products.

YOUR LOCAL NTS DISTRIBUTOR: